

Specification

Solid medium for the enumeration of heterotrophic microorganisms in treated waters according to Pharmacopoeial Method.

Presentation

	Packaging Details	Shelf Life	Storage
30 Membrane filtration plates 55 mm Plates for filtration purposes with: 9 ± 1 ml	1 box containing: 6 plastic bags with 5 plates of 55 mm/ bag.	6 months	2-25 °C

Composition

Composition (g/l):	
Proteose peptone.....	0.500
Casein hydrolysate (Tryptone).....	0.500
Yeast extract.....	0.500
D(+)-Glucose.....	0.500
Starch.....	0.500
Sodium pyruvate.....	0.300
Dipotassium hydrogen phosphate.....	0.300
Magnesium sulphate (anhydrous).....	0.024
Agar.....	15.000

Description /Technique

Description

R2A Agar was proposed in 1979 by Reasoner and Goldenreich and a few years later accepted by the APHA as an alternative medium for the enumeration of stressed cells in treated potable water. The culture medium has also been adopted by the European Pharmacopoeia for the control of purified water.

The use of nutrient rich media like PCA or TSA allows the growth of most microbes, but does not permit the recuperation of stressed or chlorine resistant organisms. Using a medium like R2A with low nutrients in combination with a lower temperature and longer incubation time it is possible to induce the resuscitation of these damaged cells.

In R2A Agar the source of nitrogen is the peptone and Yeast Extract supplies the vitamins and growth factors. The source of carbon is dextrose and magnesium sulfate and potassium phosphate maintain the osmotic pressure. The starch is a detoxifier and sodium pyruvate increases the recuperation of stressed cells. The agar acts as gelling agent.

Technique

The water sample must be processed as quickly as possible. If it is not possible to process within the first 6 hours, the sample must be refrigerated, but not for more than 30 hours.

Collect, dilute and prepare samples and volumes to be filtered as required according to specifications, directives, official standard regulations and/or expected results.

Filter the sample through a 0.45 mm pore membrane and apply it onto the surface of the agar.

After incubating at 35°C, an incubation period of 3-5 days is recommended. In most circumstances an incubation temperature of 20-25°C for 5-7 days is more effective. Plates must be protected against dehydration.

After incubation, enumerate all the colonies that have appeared onto the surface of the membrane.

Calculate total microbial count per ml of sample by multiplying the average number of colonies per plate by the inverse dilution factor.

Report results as Colony Forming Unit (CFU's) per ml along with incubation time and temperature.

Quality control**Physical/Chemical control**

Color : Pale yellow

pH: 7.2 ± 0.2 at 25°C

Microbiological control

Membrane Filtration; Practical range 10-100 CFU (productivity) according Eur. Pharm.

Aerobiosis. Incubation at 32,5°C ±2,5. Reading at 24-72 h for bacteria and 5-7 days for yeasts and moulds

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Ps. aeruginosa and *E. coli* double incubation temp. 30-35 °C / 20-25 °C

Microbiological control according to ISO 11133:2014/A1:2018; A2:2020.

Microorganism**Growth**

<i>Ps. aeruginosa</i> ATCC® 9027, WDCM 00026	Good (≥70%)
<i>Bacillus subtilis</i> ATCC® 6633, WDCM 00003	Good (≥70%)
<i>Escherichia coli</i> ATCC® 8739, WDCM 00012	Good (≥70%)
<i>Aspergillus brasiliensis</i> ATCC® 16404, WDCM 00053	Good (≥70%)
<i>Candida albicans</i> ATCC® 10231, WDCM 00054	Good (≥70%)
<i>Staphylococcus aureus</i> ATCC® 6538, WDCM 00032	Good (≥70%)
<i>E. coli</i> ATCC® 8739, WDCM 00012 (20-25°C)	Good (≥70%)
<i>Ps. aeruginosa</i> ATCC® 9027, WDCM 00026 (20-25°C)	Good (≥70%)

Sterility control

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Bibliography

- ATLAS, R.M. (1995) Handbook of Media for Environmental Microbiology. CRC Press. Boca Raton. Fla. USA.
- CLESCERI, L.S., A.E. GREENBERG and A.D. EATON (1998) Standard Methods for the Examination of Water and Wastewater. 20th ed. APHA Washington D.C. USA.
- EATON, A.D., A.E. GREENBERG and L.S. CLESCERI (1995). Standard Methods for the Examination of Water and Wastewater. 19th ed. APHA Washington D.C. USA.
- EUROPEAN PHARMACOPOEIA. 10th ed. Suppl 6.3 (2020). General Monographs. Water for injections. (pg. 4339) EDQM. Council of Europe. Strasbourg.
- GREENBERG, A.E., R.R. TRUSSELL and L.S. CLESCERI (1985). Standard Methods for the Examination of Water and Wastewater. 16th ed. APHA-AWWA-WPCF. Washington D.C. USA.
- REASONER, D.J. and E.E. GELDREICH (1979) A new Medium for the enumeration and subculture of bacteria from potable water. Abstracts of Annual Meeting. ASM 79th Meeting. Paper #N7.
- Van SOETSBERGER, A.A. and C.H. LEE (1969) Pour plates or streak plates?. Appl. Microbiol. 18:1092 -1094.