

## Specification

Solid medium for the enumeration and cultivation of fungi according to the Pharmacopeial Harmonised Method and ISO standard.

## Presentation

	Packaging Details	Shelf Life	Storage
20 Plates/Ird 90 mm - Triple Wrapping with: 21 ± 2 ml	1 box with 2 BOPP bags (triple wrapping) with 10 plates/bag. Every pack exhibits a irradiation indicator stacked on the side of the bag (8-14 KGy) with desiccant. LATERAL LABELLING	8 months	15-25 °C

## Composition

Composition (g/l):	
D(+)-Glucose.....	40.0
Peptone from casein .....	5.0
Meat Peptone.....	5.0
Agar.....	15.0

## Description /Technique

Sabouraud Dextrose Agar is a modification of the classical Sabouraud medium for the cultivation of fungi. This new formula helps to maintain the morphology of fungi, providing a reliable medium for both cultivation and differentiation. Its selectivity is due to a low pH and a high glucose concentration, which together with incubation at a relatively lower temperature (20 -25°C) favours the growth of fungi while discouraging that of bacteria. The mixture of peptones employed has been selected to provide the fungi with all their nitrogen requirements.

Spread the plate streaking methodology or by spiral method.

Each laboratory must evaluate the results according to their specifications.

**Attention:** Petri plates are used for monitoring the microbiological contamination of surface and air inside cleanrooms, isolators, RABS, food industries and hospitals. The double/triple irradiated wrapping ensures that the package itself doesn't contaminate the environment as the first wrapper is removed just before entering the clean area. Wrapping resistant to hydrogen peroxide vapors penetration.

## Quality control

### Physical/Chemical control

Color : Straw-coloured yellow      pH: 5.6 ± 0.2 at 25°C

### Microbiological control

Growth Promotion Test 50-100 CFU according to harmonized Pharmacopoeia monographs (EP) and test methods & ISO 11133:2014/A1:2018

Spiral Spreading: Practical range 50 - 100 CFU (productivity).

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Aerobiosis. Incubation at 20-25°C and 30-35°C. Reading at 72 hours for bacteria and 3-5 days for yeast and moulds.

### **Microorganism**

*Aspergillus brasiliensis* ATCC® 16404, WDCM 00053  
*S. cerevisiae* ATCC® 9763, WDCM 00058  
*Candida albicans* ATCC® 10231, WDCM 00054 (20-25°C)  
*Candida albicans* ATCC® 10231, WDCM 00054 (30-35°C)

### **Growth**

Good (≥70%)  
 Good (≥70%)  
 Good (≥70%)  
 Good (≥70%)

### Sterility control

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.  
 Check at 7 days after incubation in same conditions.

## Bibliography

- AJELLO, L. (1957) Cultural Methods for Human Pathogenic Fungi J. Chron. Dis. 5:545-551.
- COLIPA (1997) Guidelines on Microbial Quality Management (MQM). Brussels.
- EUROPEAN PHARMACOPOEIA 11.0 (2023) 11th ed § 2.6.12. Microbiological examination of Non-sterile products.: Microbial Enumeration Test. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- EUROPEAN PHARMACOPOEIA 11.0 (2023) 11th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- GEORGE, L.K., AJELLO, L. & PAPAGEORGE, C. (1954) Use of Cycloheximide in the Selective Isolation of Fungi Pathogenic to Man. J. Lab. Clin. Med, 44 (422-428).
- HANTSCHKE, D. (1968) Mykosen, 11, (769-778).
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 4973:2023. Quality control of culture media and diluents used in cosmetics standards
- ISO 11930:2019. Evaluation of the antimicrobial protection of a cosmetic product.
- ISO 16212 Standard (2017) Cosmetics - Microbiology - Enumeration of yeast and mould.
- PAGANO, J. LEVIN, J.D. and TREJO, W. (1957-58) Diagnostic Medium for Differentiation of Species of Candida. Antibiotics Annual, 137-143.
- SABOURAUD, R. (1910) Les Teignes. Masson, Paris.
- USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA