

Specification

A sterile selective supplement used for isolation and presumptive identification of *Clostridium perfringens*, according to ISO Standards and other regulations.

Presentation

	Packaging Details	Shelf Life	Storage
10 Freeze dried vials Vial with: 3 ± 0,1 g	22±0,25 x 55±0,5 mm glass vials, tag labelled, White plastic cap - 10 vials per box.	49 months	2-25°C

Composition

Composition (g/vial)

D-Cycloserine.....0.100

Reconstitute the original freeze-dried vial
by adding
Sterile Distilled Water.....6 ml

NOTE: Each vial is sufficient to supplement 250
ml of medium Base: TSC Agar Base.

Description /Technique

Description:

D-cycloserine selective supplement is added to TSC Agar in order to obtain a final selective medium which has the advantage to simplify the counting of plates with high numbers of colonies because smaller colonies of *C. perfringens* are formed. Sodium metabisulphite and ferric ammonium citrate are used as an indicator of sulphite reduction made by *Clostridium perfringens* spp. that produce black colonies in TSC agar.

Technique:

Collect, dilute and prepare samples and volumes as required according to specifications, directives, official standard regulations and/or expected results.

Reconstitute the vial with 6ml of sterile diluent in aseptic conditions and add it to 250 ml of melted Agar base cooled to 50°C.

Do not overheat once supplemented.

Pour the complete medium into Petri dishes (or tubes) and, once solidified on a flat surface, spread the plates either by streaking by spiral method or dilution banc.

Incubate the plates in anaerobic atmosphere at 35 ± 2°C for 20-24h. To obtain a more selective medium, incubated at 44° C ± 1.

Incubation times longer than those mentioned above or different incubation temperatures may be required depending on the sample or the specifications.

After incubation, count all the colonies that have appeared onto the surface of the agar.

C. perfringens grows in black colonies, due to the iron sulfide precipitation.

Presumptive isolation of *Clostridium perfringens* must be confirmed by further microbiological and biochemical tests.

Quality control

Physical/Chemical control

Color : White-Gray

pH: at 25°C

Microbiological control

Add 1 vial to 225/ 250 ml of medium base. DO NOT HEAT once supplemente

Reconstitute 1 vial as indicated in COMPOSITION; shake and dissolve completely

Anaerobiosis. Incubation at 44 ± 1 °C during 21 ± 3h.

Microorganism

Clostridium perfringens ATCC® 13124, WDCM 00007, NCTC® 8237

Clostridium perfringens ATCC® 10543, WDCM 00174

Bacillus subtilis ATCC® 6633, WDCM 00003

Growth

Good - black colonies

Good - black colonies

Inhibited

Sterility Control

Add 5ml of the sample to 100ml of TSB and to 100ml Thioglycollate

Incubation 48 hours at 30-35°C and 48 hours at 20-25°C: NO GROWTH

Check at 7 days after incubation in same conditions

Bibliography

- ATLAS, R.M., LC. PARKS (1993) Handbook of Microbiological Media. CRC Press, Inc. London.
- DIN Standard 10165. Referenz Verfahren für Bestimmung von *Clostridium perfringens*. Fleisch und Fleischerzeugnissen.
- DOWNES, F.P. & K. ITO (2001) Compendium of Methods for the Microbiological Examination of Foods. 4th ed. American Public Health Association. Washington.
- DIRECTIVA 2015/1787/UE de la Comisión por la que se modifica la Directiva 98/ 83/CE relativa a la calidad de las aguas destinadas al consumo humano (DO L260 de 7.10.2015 pg 6 y ss)
- FDA (Food and Drug Administrations) (1998) Bacteriological Analytical Manual. 8th ed. Revision A. AOAC International Inc. Gaithersburg. MD.
- ISO 7937 (2004) Microbiology of Food and Animal Feeding Stuffs. Horizontal Method for Enumeration of *C. perfringens*. Colony-count technique.
- ISO Norma 6461-2 (1986) Water Quality.- Detection and enumeration of the spores of sulfite-reducing anaerobes (Clostridia).- Part 2: Method by Membrane Filtration.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 14189 (2013) Water quality. Enumeration of *Clostridium perfringens* — Method using membrane filtration
- SMITH, L.D. (1981) Clostridial Anaerobic Infections, in Diagnostic Procedures for Bacterial Mycotic and Parasitic Infections. 6th ed. APHA. Washington.
- UNE-EN ISO 11133 (2014). Microbiología de los alimentos para consumo humano, alimentación animal y agua.-Preparación, producción, conservación y ensayos de rendimiento de los medios de cultivo.